

Come to *Chesapeake Plastic Surgery & Aesthetics* for all of your dermal filler needs!

Rebate Incentive

The Restylane® family of products is offering a special rebate incentive. For a limited time only, you can receive a \$50 rebate on one 1-mL syringe or a \$100 rebate on two 1-mL syringes. Just simply fill out the Restylane® Rebate Form when you come in. **Offer expires October 31st, 2008.**

Contact Jessica at (410) 646-3226 or jclow@4cpsa.com to schedule your appointment today!

This fall make the ultimate fashion statement.

Nasolabial folds before After one Restylane® 1 mL syringe After two Restylane® 1 mL syringes
Individual results may vary. Individual results may vary.

Restylane®
redefining beauty™

Perlane®

Important Safety Considerations of Restylane®

Restylane® restores volume and fullness to the skin to correct moderate to severe facial wrinkles and folds, such as the lines from your nose to the corners of your mouth (nasolabial folds). After your treatment, you might have some swelling, redness, pain, bruising, and tenderness. This will normally last less than seven days. Although rare, red or swollen small bumps may occur. If you have had facial cold sores before, an injection can cause another outbreak. In rare circumstances, the doctor may inject into a blood vessel, which can damage the skin. To avoid bruising and bleeding, you should not use Restylane® if you have recently used drugs that thin your blood or prevent clotting. If you are pregnant, breastfeeding, or under 18, you should not use Restylane®.

Restylane® should not be used by people with previous bad allergies, particularly to certain microorganisms known as gram positive bacteria, or by people with previous bad allergies to drugs that have required in-hospital treatment. Restylane® should not be injected anywhere except the skin or just under the skin.

The use of Restylane® at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. Use of Restylane® in these instances could delay healing or make your skin problems worse.

Restylane® and Perlane® are available only through a licensed practitioner. Complete product and safety information for both products is available at www.RestylaneUSA.com. To opt out of future e-mails contact the physician's office.

Important Safety Considerations of Perlane®

Perlane® restores volume and fullness to the skin to correct moderate to severe facial wrinkles and folds, such as the lines from your nose to the corners of your mouth (nasolabial folds). After your treatment, you might have some swelling, redness, pain, bruising, and tenderness. This will normally last less than seven days. Although rare, red or swollen small bumps may occur. If you have had facial cold sores before, an injection can cause another outbreak. In rare circumstances, the doctor may inject into a blood vessel, which can damage the skin. To avoid bruising and bleeding, you should not use Perlane® if you have recently used drugs that thin your blood or prevent clotting. If you are pregnant, breastfeeding, or under 18, you should not use Perlane®.

Perlane® should not be used by people with previous bad allergies, particularly to certain microorganisms known as gram positive bacteria; by people with previous bad allergies to drugs that have required in-hospital treatment; or by people with bleeding disorders. Perlane® should not be injected anywhere except the skin or just under the skin.

The use of Perlane® at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. Use of Perlane® in these instances could delay healing or make your skin problems worse.